

# Interferon Treatment of Chronic Active Hepatitis C During Therapy of Acute Lymphoblastic Leukemia

Peter Waldron

Department of Pediatrics, University of Virginia, Charlottesville, Virginia

Due to concerns that antineoplastic therapy produces prolonged decrease in immune function, interferon treatment of chronic active hepatitis C (CAHC) has been used only at one year or longer after the end of cancer therapy. We report the experience of an 11-year-old who developed symptomatic CAHC at the start of maintenance therapy for testicular relapse of acute lymphoblastic leukemia (ALL). Significant dose reduction of maintenance therapy did not improve the tolerance of antileukemic treatment. In an effort to improve his liver disease and to deliver effective antileukemic therapy, interferon alpha and an alternative maintenance therapy regimen for ALL were initiated. The patient tolerated the combined therapy well. Interferon therapy was continued for 27 months, which was three months from the end of antineoplastic therapy. At that time serum transaminase values were normal, and no HCV viral genome was detectable. Viral genome was detected 10 months later. The combined effects of interferon and antineoplastic therapy resulted in myelosuppression requiring dose reduction of both treatments. The patient remains asymptomatic and with no evidence of recurrent leukemia more than six years from diagnosis of relapse. The effect on the status of this patient's CAHC was similar to that reported among leukemic patients who underwent an interferon course more than one year from the end of antineoplastic therapy. Interferon treatment of CAHC can be given concomitantly with antineoplastic therapy. *Am. J. Hematol.* 61:130–134, 1999.

© 1999 Wiley-Liss, Inc.

**Key words:** hepatitis C; interferon; acute lymphoblastic leukemia

## INTRODUCTION

Hepatitis C virus (HCV) is the etiologic agent of as much as 90% of posttransfusion hepatitis prior to 1990 [1,2]. Individuals with acute lymphoblastic leukemia (ALL) are at risk for HCV infection since they usually receive at least one, and typically multiple, transfusions of blood products during the course of their disease [3]. The frequency of antibody to HCV in children with leukemia ranges from 1 to 43% [4–7]. Reports of interferon treatment of chronic active viral hepatitis have stressed the importance of waiting as long as two years after the completion of antineoplastic therapy to ensure immune competence [8,9]. There are no prior reports of the use of interferon- $\alpha$  concomitant with cytotoxic antileukemic therapy.

We report on a patient who had jaundice, malaise, antibody to HCV, and the histologic appearance of chronic active hepatitis at the time of testicular relapse of ALL. During systemic therapy for testicular relapse, he developed transaminase elevation to greater than 15

times normal and total bilirubin (TB) greater than 6.0 mg/dl. Recombinant interferon- $\alpha$ 2a (Ifn- $\alpha$ ) was initiated to treat his liver disease and to allow effective antileukemic therapy to be delivered. The patient tolerated Ifn- $\alpha$  therapy well, and completed systemic treatment for testicular relapse of ALL.

## CASE REPORT

In October 1988 the patient was eight years old when he was diagnosed with B-cell precursor ALL. He was treated according to a contemporary protocol (Pediatric Oncology Group #8602 regimen C). During the first 20 weeks of therapy he received eight units of packed red

Correspondence to: Peter Waldron, M.D., Department of Pediatrics, Box 386 HSC, University of Virginia, Charlottesville, VA 22908; E-mail [pew6f@virginia.edu](mailto:pew6f@virginia.edu)


Received for publication 16 July 1998; Accepted 3 February 1999

blood cells and 58 units of random donor platelets. Symptoms of hepatitis were not manifest until week 137 of treatment when he developed prolonged malaise, lethargy, vomiting, and diarrhea. At that time his alanine amino transferase (ALT) / aspartate aminotransferase (AST) were 130/124 and his TB was 4.8 mg/dl. Antibody to hepatitis A virus, hepatitis B surface antigen, and core protein were not detected, nor was hepatitis B surface antigen. HCV serology using a first generation ELISA assay was positive. The treating physician chose to stop maintenance at week 142 of a planned 156 week course due to the serologic diagnosis of HCV, the symptoms of hepatitis, and poor tolerance of maintenance leukemic therapy. Off therapy evaluation showed no evidence of leukemia in bone marrow or cerebrospinal fluid (CSF) and a normal physical exam.

One month after stopping therapy, a routine physical exam showed that both testes were increased in size and hard. Testicular biopsy showed a leukemic infiltrate. During the same anesthesia a liver biopsy was done. This showed hepatocellular necrosis with grade 2 periportal inflammation, and stage 2 portal fibrosis with focal areas of stage 3 bridging fibrosis [10]. Repeat bone marrow and CSF exam were normal. The patient underwent 2500 cGy radiation therapy to the testes and standard four-drug systemic reinduction for ALL. Consolidation consisted of cyclophosphamide 1 gm/M<sup>2</sup>/dose on days 1 and 14, daily oral 6-mercaptopurine (6-MP) 50 mg/M<sup>2</sup>, and four-day courses starting days 1, 8, 15, and 22 of cytosine arabinoside (Ara-C) 75 mg/M<sup>2</sup>/d IV. A week prior to liver biopsy ALT/AST were 293/192. One week after liver biopsy, at the start of induction therapy, ALT/AST/TB were 925/736/1.0. At the end of induction or start of consolidation ALT/AST/TB were 447/175/0.8.

Planned maintenance therapy was daily oral 6-MP (50 mg/M<sup>2</sup>); weekly oral methotrexate (MTX) 20 mg/M<sup>2</sup>, and every four weeks vincristine and five-day prednisone 40 mg/M<sup>2</sup>/d. Central nervous system prophylaxis was with intrathecal MTX and oral MTX was not given during that week. At the beginning of maintenance ALT/AST/TB were 125/89/0.6. MTX was given IT during week one of maintenance therapy, and the first oral MTX was given at week two. At the start of week three ALT/AST/TB were 101/90/3.6, and the patient complained of mild to moderate malaise. 6-MP and MTX were held due to elevated TB. One week later AST/ALT/TB were 289/178/1.7, and oral medication was resumed at 25% and 17% of the planned 6-MP and MTX dose, respectively. After one week of this dose schedule AST/ALT/TB were 984/490/1.2.

Due to the need for systemic ALL therapy and the poor tolerance of even small fractions of the intended doses of 6-MP and MTX, an alternative therapeutic approach was pursued. Ifn- $\alpha$  therapy was initiated at a dose of 1 million units (MU) subcutaneously (0.66 MU/M<sup>2</sup>) three days per



**Fig. 1.** Patient's serum alanine amino transferase (ALT) represented as actual value/upper limit of institutional normal range or times normal (x's NI); serum total bilirubin (TB) represented as actual values, and interferon (Ifn) dose per week in million units (MU).

week. This dose was increased with one intermediary step to 2.0 MU per dose (1.32 MU/M<sup>2</sup>/dose) three days per week (see Figure 1). Maintenance antileukemic therapy was altered to decrease the exposure to hepatotoxic agents. It utilized noncross resistant pairs of drugs and consisted of etoposide 150 mg/M<sup>2</sup>/dose IV, Ara-C 300 mg/M<sup>2</sup>/dose IV alternating every four weeks with cyclophosphamide 600 mg/M<sup>2</sup>/dose, vincristine 1.5 mg/M<sup>2</sup>/dose and five-day prednisone 40 mg/M<sup>2</sup>/day [11]. Secondary to prolonged neutropenia and thrombocytopenia the standard doses of 150 mg/M<sup>2</sup> of etoposide and 300 mg/M<sup>2</sup> of Ara-C were reduced to 100 mg/M<sup>2</sup>/dose and 150 mg/M<sup>2</sup>/dose, respectively. Vincristine was not given when total TB was greater than 1.5 mg/dl. Ifn- $\alpha$  dose was adjusted to maintain platelets greater than 75,000/mm<sup>3</sup> and neutrophils greater than 1,000/mm<sup>3</sup>. Ultimately, this resulted in a dose of Ifn- $\alpha$  of 1.5 MU two days per week. From month 7 to month 27 the Ifn- $\alpha$  dose per M<sup>2</sup> per week varied from 1.9 to 1.74 MU due to delivery of a constant dose with growth in body surface area. Maintenance leukemia therapy was continued for a total of 24 months. The patient continued to receive Ifn- $\alpha$  therapy until one month after his ALT had decreased to within the normal range, at which time reverse transcriptase-polymerase chain reaction (RT-PCR) for HCV was negative. This was three months after the end of cytotoxic leukemia therapy resulting in a total of 27 months of Ifn- $\alpha$  therapy. Repeat RT-PCR for HCV was positive at months 37 and 53. Androgen replacement therapy was begun after completion of the antileukemic therapy. The patient remains in remission 72 months from testicular relapse. Current clinical status including activity level and growth is normal. His peripheral blood counts are normal with the exception of a platelet count between 90,000 and 140,000/mm<sup>3</sup>. A bone marrow exam one year from the end of leukemia therapy and nine months from

the end of interferon therapy showed normal cellularity with slightly decreased megakaryocyte number.

## DISCUSSION

We have described the clinical course of a patient with ALL who was diagnosed with chronic active hepatitis C (CAHC) simultaneous with diagnosis of an early, isolated, testicular relapse of ALL. Although ALL induction and consolidation therapy were well tolerated he showed clinical and laboratory signs of exacerbation of CAHC during the maintenance phase of therapy. Ifn- $\alpha$  and alternative ALL maintenance therapy were used simultaneously. Low platelet and neutrophil number required dose alteration for both Ifn- $\alpha$  and cytotoxic chemotherapy, but the treatments were otherwise well tolerated. This is the first reported experience with simultaneous administration of cytotoxic therapy for ALL and Ifn- $\alpha$  for CAHC.

HCV infection is rare in the pediatric age group. Two large studies found similar frequencies of 0.05 and 0.1% of antibody to HCV among general adolescent populations in the United States [12,13]. Individuals who receive blood products, including those with malignancies [14,15], hemophilia [16,17], sickle cell disease [18], and thalassemia [19,20] have a higher incidence of antibody to HCV compared with the general population. The impact of HCV infection on the tolerance of simultaneous therapy and the natural history of these diseases is only beginning to be defined. A retrospective study of 87 southern Italian children at the end of their treatment for ALL found that 32% were HCV antibody positive. Those with antibody to HCV had more frequent elevation of ALT and TB and had more frequent interruption of maintenance ALL therapy (82% versus <2%) compared with the ALL patients who were HCV antibody negative. Five of the 28 patients with HCV antibody developed signs of chronic liver disease [7]. The frequency of HCV infection among these children from southern Italy is similar to the 43% reported among 102 children with ALL from northern Italy [6]. However, it is in marked contrast to the finding of 1 of 98 children with hematologic malignancies from central England who had anti-HCV antibody or had detectable HCV by RT-PCR [4] and the 13% of 24 children with acute leukemia who were antibody positive against HCV from Massachusetts [14].

Ifn- $\alpha$  was established as a therapy that was effective for some adult patients with chronic active non-A, non-B hepatitis prior to the identification of the HCV [21,22]. Since then the experience with more than 1,800 adult patients has been reported, allowing clearer recommendations for dose schedule and selection of patients [23–25]. The use of Ifn- $\alpha$  in children with CAHC was first reported by Ruiz-Moreno et al., who treated subjects who had no underlying disease with 3 MU/M<sup>2</sup> Ifn- $\alpha$  three

times per week for six months. Five of the 11 evaluable patients had normal values for ALT and eight were negative for HCV RNA at 24 months from the start of treatment. All children were reported to have an improvement in their liver histology [26]. The optimal dose, timing, and strategy for use of Ifn- $\alpha$  in children with HCV infection is not yet defined. Eight additional studies of Ifn- $\alpha$  for children with CAHC have been reported [8,27–33]. The frequency of sustained complete response defined as absence of detectable viral genome and normal ALT typically at 12 months after the end of treatment has consistently been in the range of 30 to 40%. The most relevant experience to our patient is that of Komatsu et al., who treated 13 individuals, who had completed leukemia therapy, with Ifn- $\alpha$  for CAHC. All subjects were at least two years from the end of leukemia therapy due to a concern that the cytotoxic chemotherapy may cause a prolonged immunosuppressed state and that immunocompetence was necessary for Ifn- $\alpha$  efficacy. These children were given 1 MU per kg to a maximum of 6 MU daily for two weeks and then three times per week for an additional 22 weeks. Prior to Ifn- $\alpha$  therapy all children had detectable HCV genome by RT-PCR and had ALT above normal. Twelve months after completion of Ifn- $\alpha$  therapy four of 13 were negative for the viral genome and eight of 13, including all RT-PCR negative patients, had normal ALT values. Complications of Ifn- $\alpha$  therapy included an influenza-like syndrome in all children that generally lasted less than 10 days; slight, transient hair loss in two children and a “small” decrease in leukocyte and platelet count in all children, that did not necessitate dosage reduction [8].

Acute adverse effects of treatment for our patient were mild malaise and myalgia for four weeks duration, grade two thrombocytopenia and grade three and four neutropenia. However, the assessment of the myelosuppressive effect of Ifn- $\alpha$  is confounded by the simultaneous administration of cytotoxic therapy. He also has chronic grade 1 thrombocytopenia. The mechanism of myelosuppression may be a direct effect of Ifn- $\alpha$  on bone marrow progenitor cells [34]. In one series of patients with CAHC treated with Ifn- $\alpha$  the reported frequency of grade 3 thrombocytopenia was 10% and grade 3 neutropenia was 20% [35]. Ultimately the tolerated dose of Ifn- $\alpha$  for our patient in the context of leukemic therapy was 1.5 MU two days per week (mean dose of 1.8 MU/M<sup>2</sup> per week), which was 20% of a recommended dose of 9 MU/M<sup>2</sup> per week. At this interferon dose it was possible to deliver 67% of planned leukemic maintenance therapy. Thus, the simultaneous treatment of hepatitis and leukemia did compromise the drug delivery for the individual diseases. Additionally, Ifn- $\alpha$  may have had some antileukemic effect in view of clinical and in vitro data indicating efficacy of Ifn- $\alpha$  against ALL [36–38]. The strategy employed to provide an adequate treatment

of hepatitis was to continue the drug to a biological end-point of normalization of ALT value. This resulted in a prolonged (27 months), but low-dose course. The total dose delivered was 240 MU/M<sup>2</sup>, which is similar to the total dose used by Ruiz-Monero et al. [26] and Komatsu et al. [8] but distributed over a longer time.

Dose intensity, defined as the amount of drug delivered per period of time, may be significant when considering patients who develop CAHC during treatment for ALL. Dibenedetto et al. described an 80% frequency of suspension of ALL maintenance therapy in the 29 patients who were HCV antibody positive, but only a 2% frequency of suspensions in those who were HCV antibody negative [7]. Thus, for these patients ALL therapy dose intensity was decreased when hepatitis C infection was present. An early randomized study by Pinkel et al. found that children with ALL who received full-dose of maintenance therapy had a longer median duration of complete response and greater likelihood of complete remission at the end of therapy than a group who were treated with half-dose maintenance therapy [39]. Recently both Italian and German investigators reported a greater event-free survival among ALL patients who received a higher portion of the intended maintenance therapy [40,41]. Potentially, early recognition and treatment with Ifn- $\alpha$  of acute HCV could allow delivery of a greater dose intensity of antileukemia therapy.

The experience with this patient demonstrates that Ifn- $\alpha$  can be given simultaneously with leukemic therapy. Our single patient had an efficacy and tolerance to Ifn- $\alpha$  that is within the range of findings reported for subjects who were not receiving antineoplastic therapy. The simultaneous administration of Ifn- $\alpha$  and antineoplastic therapy may allow the delivery of a greater proportion of planned therapy, with greater dose intensity and a decreased frequency of disease recurrence.

## Acknowledgments

The author thanks Drs. J. Sutphen and S. Borowitz for helpful discussions of clinical management, Drs. M. Lovell and J. Iezzoni for interpretation of the liver biopsy material and Dr. P deAlarcon for critique of the manuscript.

## REFERENCES

1. Alter HJ. To C or not to C: these are the questions. *Blood* 1995;85:1681-1695.
2. Aach RD, Stevens CE, Hollinger FB, Mosley JW, Peterson DA, Taylor PE, Johnson RG, Barbosa LH, Nemo G. Hepatitis C virus infection in post-transfusion hepatitis. *N Engl J Med* 1991;325:1325-1329.
3. Hetherington ML, Buchanan GR. Elevated serum transaminase values during therapy for acute lymphoblastic leukemia correlate with prior blood transfusions. *Cancer* 1988;62:1614-1618.
4. Myers B, Irving W, Hollingsworth R, Readett D, Lilleyman JS, Dolan G. Hepatitis C virus infection in multi-transfused children with haematological malignancy. *Br J Haematol* 1995;91:480-482.
5. Arico M, Maggiore G, Silini E, Bono F, Vigano C, Cerino A, Mondelli MU. Hepatitis C virus infection in children treated for acute lymphoblastic leukemia. *Blood* 1994;84:2919-2922.
6. Locasciulli A, Cavalletto D, Pontisso P. Hepatitis C virus serum markers and liver disease in children with leukemia during and after chemotherapy. *Blood* 1993;82:2564-2567.
7. Dibenedetto SP, Ragusa R, Sciacca A, Di Cataldo A, Miraglia V, D'Amico S, Lo Nigro L, Ippolito AM. Incidence and morbidity of infection by hepatitis C virus in children with acute lymphoblastic leukaemia. *Eur J Pediatr* 1994;153:271-275.
8. Komatsu H, Fujisawa T, Inui A, Miyakawa Y, Onoue M, Sekine I, Hanada R, Yamamoto K. Efficacy of interferon in treating chronic hepatitis C in children with a history of acute leukemia. *Blood* 1996;87:4072-4075.
9. Cetingul N, Kavakli K, Vergin C, Oztup S, Nisli G, Aydogdu S, Ozcar T, Bilgic A, Yuce G, Yagci R. Alpha-interferon-2b treatment for chronic hepatitis-B infection in children with cancer. *J Trop Pediatr* 1996;42:263-266.
10. Batts KP, Ludwig J. Chronic hepatitis: an update on terminology and reporting. *Am J Surg Pathol* 1995;19:1409-1417.
11. Rivera GK, Buchanan G, Boyett JM, Camitta B, Ochs J, Kalwinsky D, Amylon M, Vietti TJ, Crist WM. Intensive retreatment of childhood acute lymphoblastic leukemia in first bone marrow relapse: a pediatric oncology group study. *N Engl J Med* 1986;315:273-278.
12. Murphy EL, Bryzman S, Williams AE, Co-Chien H, Schrieber GB, Ownby HE, Gilcher RO, Kleinman SH, Matijas L, Thomson RA, Nemo GJ. Demographic determinants of hepatitis C virus seroprevalence among blood donors. *J Am Med Assoc* 1996;275:995-1000.
13. Jonas MM, Robertson LM, Middleman AB. Low prevalence of antibody to hepatitis C virus in an urban adolescent population. *J Pediatr* 1997;131:314-316.
14. Monteleone PM, Andrzejewski C, Kelleher JF. Prevalence of antibodies to hepatitis C virus in transfused children with cancer. *Am J Pediatr Hematol Oncol* 1994;16:309-313.
15. Cesaro S, Petris MG, Rossetti F, Cusinato R, Pipan C, Guido M, Masiero L, Botta GA, Meloni GA, Zanenco L. Chronic hepatitis C virus infection after treatment for pediatric malignancy. *Blood* 1997;90:1315-1320.
16. Kanesaki T, Kinoshita S, Tsujino G, Yoshioka K, Ikegami N. Hepatitis C virus infection in children with hemophilia: characterization of antibody response to four different antigens and relationship of antibody response, viremia, and hepatic dysfunction. *J Pediatr* 1993;123:381-387.
17. Wagner N, Rothauwe HW. Hepatitis C contributes to liver disease in children and adolescents with hemophilia. *Klin Pädiatr* 1994;206:40-44.
18. Hasan MF, Marsh F, Posner G, Bellevue R, Dosik H, Suatengco R, Ramani N. Chronic hepatitis C in patients with sickle cell disease. *Am J Gastroenterol* 1996;91:1204-1206.
19. Ni YH, Chang MH, Lin KH, Chen PJ, Lin DT, Hsu HY, Chen DS. Hepatitis C viral infection in thalassemic children: Clinical and molecular studies. *Pediatr Res* 1996;39:323-328.
20. Angelucci E. Antibodies to hepatitis C virus in thalassemia. *Haematologica* 1994;79:353-355.
21. Jacyna MR, Brooks MG, Loke RH, Main J, Murray-Lyon IM, Thomas HC. Randomised controlled trial of interferon alfa (lymphoblastoid interferon) in chronic non-A, non-B hepatitis. *Br Med J* 1989;298:80-82.
22. Schvarcz R, Weiland O, Wejstal R, Norkrans G, Fryden A, Foberg U. A randomised controlled open study of interferon alpha-2b treatment of chronic non-A, non-B post-transfusion hepatitis: no correlation of outcome to presence of hepatitis C virus antibodies. *Scand J Infect Dis* 1989; 21:617-625.
23. Poynard T, Leroy V, Cohard M, Thevenot T, Mathurin P, Polon P,



- Zarski JP. Meta-analysis of interferon randomized trials in the treatment of viral hepatitis C: effects of dose and duration. *Hepatology* 1996;24:778–789.
24. Powell DW. NIH Consensus Development Panel: management of hepatitis C. *Hepatology* 1997;26(Suppl 1):2S–9S.
25. Lindsay KL. Therapy of hepatitis C: overview. *Hepatology* 1997;26(3 Suppl 1):71S–77S.
26. Ruiz-Monero M, Rua MJ, Castillo I, Garcia-Novo MD, Santos M, Navas S, Carreno V. Treatment of children with chronic hepatitis C with recombinant interferon-alpha: a pilot study. *Hepatology* 1992;16:882–885.
27. Matsuoka S, Mori K, Nakano O, Yuasa Y, Taguchi Y, Hayabuchi Y, Kuroda Y. Efficacy of interferons in treating children with chronic hepatitis C. *Eur J Pediatr* 1997;156:704–708.
28. Bortolotti F, Giacchino R, Vajro P, Barbera C, Crivellaro C, Alberti A, Nebbia G, Zancan L, De Moliner L, Bertolini A, Balli F, Callea F. Recombinant interferon-alfa therapy in children with chronic hepatitis C. *Hepatology* 1995;22:1623–1627.
29. Marcellini M, Kondili L, Comparcola D, Spada E, Sartorelli, Palumbo M, Rapicetta M. High dosage alpha-interferon for treatment of children and young adults with chronic hepatitis C disease. *Pediatr Infect Dis J* 1997;16:1049–1053.
30. Iorio C, Pensati R, Porzio S, Fariello I, Guida S, Vegnente A. Lymphoblastoid interferon alfa treatment in chronic hepatitis C. *Arch Dis Child* 1996;74:152–156.
31. Clemente MG, Congia M, Lai ME, Lilliu F, Lampis R, Frau F, Frau MR, Faa G, Diana G, Dessì C, Melis A, Mazzoleni AP, Cornacchia G, Cao A, De Virgiliis. Effect of iron overload on the response to recombinant interferon-alfa treatment in transfusion-dependent patients with thalassemia major and chronic hepatitis C. *J Pediatr* 1994;125:123–128.
32. Di Marco V, Lo Iacono O, Almasio P, Ciaccio C, Capra M, Rizzo M, Malizia R, Maggio A, Fabiano C, Barbaria F, Craxi A. Long-term efficacy of alpha-interferon in beta-thalassemics with chronic hepatitis C. *Blood* 1997;90:2207–2212.
33. Fujisawa T, Inui A, Ohkawa T, Miyakawa Y, Onoue M, Sekine I, Hanada R, Yamamoto K. Response to interferon therapy in children with chronic hepatitis C. *J Pediatr* 1995;127:660–662.
34. Geissler RG, Ottman OG, Kojouharoff G, Reutzel P, Eder M, Hoelzer D, Ganser A. Influence of human recombinant interferon-alpha and interferon-gamma on bone marrow progenitor cells of HIV-positive individuals. *AIDS Res Hum Retroviruses* 1992;8:521–525.
35. Poynard T, Bedossa P, Chevallier M, Mathurin P, Lemonnier C, Trepo C, Couzigou P, Payen JL, Sajus M, Costa JM. A comparison of three alfa-2b regimens for the long-term treatment of chronic non-A, non-B hepatitis. *N Engl J Med* 1995;332:1457–1462.
36. Manabe A, Yi T, Kumagai M, Campana D. Use of stroma-supported cultures of leukemic cells to assess antileukemic drugs. I. Cytotoxicity of interferon alpha in acute lymphoblastic leukemia. *Leukemia* 1993;7:1990–1995.
37. Lauer SJ, Ochs J, Pollock BH, Buchanan GR. Recombinant alpha-2B interferon treatment for childhood lymphoblastic disease in relapse. A Pediatric Oncology Group phase II study. *Cancer* 1994;74:197–202.
38. Ochs J, Brecher ML, Mahoney D, Vega R, Pollock BH, Buchanan GR, Whitehead VM, Ravindranath Y, Freeman AI. Recombinant interferon alfa given before and in combination with standard chemotherapy in children with acute lymphoblastic leukemia in first marrow relapse: a Pediatric Oncology Group pilot study. *J Clin Oncol* 1991;9:777–782.
39. Pinkel D, Hernandez K, Borella L, Holton C, Aur R, Samoy G, Pratt C. Drug dose and remission duration in childhood lymphocytic leukemia. *Cancer* 1971;27:247–256.
40. Gobrecht O, Göbel U, Graubner U, Gutjahr P, Schock, V, Spaar HJ, Janka-Schaub GE. Influence of individual dose intensive therapy on prognosis in childhood acute lymphoblastic leukemia. *Klin Pädiatr* 1992;204:230–235.
41. Dibenedetto SP, Guardabasso V, Ragusa R, Di Cataldo A, Miraglia V, D'Amico S, Ippolito AM. 6-Mercaptopurine cumulative dose: a critical factor of maintenance therapy in average risk childhood acute lymphoblastic leukemia. *Pediatr Hematol Oncol* 1994;11:251–258.